A synopsis report

On

Formulation & Evaluation Nanoemulsion for the Treatment of Topical Fungal Infection

Submitted to

IIMT University in partial fulfilment of the requirements for the award of the Degree of

DOCTOR OF PHILOSOPHY

In

Pharmaceutical Science

By

Manikant Jayner

(Roll No -2165004007)

Under the supervision of

Dr. Rustam Ekbbal

(M. Pharm. Ph.D.)

Associate Professor department of Pharmacy

Guide

IIMT COLLEGE OF MEDICAL SCIENCES

IIMT UNIVERSITY,

MEERUT, UTTAR PRADESH, INDIA.

JUNE, 2023

INDEX

S. No.	PARTCULARS	PAGE No.
1.	Introduction	6-16
2.	Review of Literature	17
3.	Aim & Objectives	18
4.	Plan of Work	19
7.	References	20

INTRODUCTION

1.1 Topical Drug Delivery System:

Drug delivery systems are procedures that are used to make sure that pharmaceuticals enter the body and reach the desired location. These frameworks should take into account a variety of requirements, from ease of delivery to suitability of treatments. A few organisations develop drug delivery techniques, market these products to drug organisations, and help other drug organisations build their own frameworks. Many of these tactics are restricted and protected. At the point when a medication is endorsed, the measurements should be exactly determined for body to ingest it. This calls for a medication delivery system that takes precise dosing into account. Drug delivery systems must also take into account how a drug is metabolised by the body. For instance, some drugs can't interact with body in this way because they are destroyed in the digestive system. Others may be extremely risky, & because of this, a period discharge strategy should be used to deliver medication for patient wellbeing. Skin drug delivery systems involve interacting a substance with the skin in a way that allows for consumption. Skin drug delivery systems, such as skin patches, are an example. Splashes applied to the nasal bodily fluid layers, inhalation vapour sprayers, eye drops, and lotions that can be scoured into the skin are only a few examples of different frameworks. These frameworks are interesting because they are typically very easy for patients to use. The objective of a medication conveyance circumstance is dependably to convey suitable dose to legitimate areas. Patients favor straightforward, simple techniques, which is reason numerous drugs are accessible as skin & enteral methodology that can be ingested or applied straightforwardly to skin. Because these procedures take greater consideration of communication over how & when the prescriptions are used, perenteral courses can be increasingly common in clinical settings, particularly for controlled substances [1, 2, 3, 4].

1.2 Topical Route of Drug Administration:

Although many substances do infiltrate skin somewhat, at generally slow rates, even though the unblemished skin is significantly less porous than other tissues, entrance of the medications & different substances through skin relies upon physiochemical properties of enter, state of skin & thought of vehicle. Drugs applied topically, basically for neighborhood activity, incorporate enemy of septic, hostile to contagious, mitigating specialists as well as skin emollients for defensive impacts. While this course can likewise be utilized for fundamental medication conveyance. Even while sweat glands, sebaceous glands, and hair follicles can all help medications given topically disperse through the skin, stratum corneum penetration is the preferred route and moves extremely slowly.

1.3 Nano-emulsion:

Nan emulsion is assessed to be a functioning and stable fluid scattering of two non-miscible fluid stages like water stage and oil stage. At the point when particles of two fluid stages alluringly associate with one another, a between facial pressure exist all over. To limit this between facial strain sur-factants are added. A limited size scattering is seen when the globule size of nano-emulsion drops especially in scope of 20-200nm. The fundamental contrast between emulsion & Nano-emulsion is that nanoemulsions are hazy & clear while emulsions are shady & coarse. What makes drug conveyance most encouraging is its simple

planning, life of soundness, among high dissolvable. Such countless applications are started in oral medications to build their ability to break down & bio-availability. Lately there is rush in examination of nano-emulsions for Transdermal use. The examinations are additionally being held for viable application in various purposes like oral, perenteral, vaginal & aspiratory application [5, 6].

A **B**

Fig1.1: A. Micro-emulsion & B. Nanoemulsion

1.3.1 Advantages:

- Because of small globules no arrangement of velvety layer & sedimentation during capacity.
- Dynamic elements of huge surface of nano-emulsion quickly enter through skin.
- Medications can be controlled through different courses like intravenous, oral or outer use.
- Hydrophilic & lipophilic can be conveyed by comparable nano-emulsion.
- Expanded patient similarity.
- Nano-emulsions give alleviating to skin & patient won't feel any of its thickness in view of its liquidity.

1.3.2 Disadvantages:

- Nano-emulsions have exceptionally low dissolvability along these lines confronting trouble on high liquefying substances.
- The instrument of creation of exact moment globules and the investment of surfactant and co surfactant is truly hard to understand.
- The exhibitions are upset because of over utilization of surfactant and co surfactant.
- Significant expense gear are required for the development of nano-emulsions [7].

1.6 Contagious Contamination:

Mycosis, another name for a skin illness caused by a parasite, is an infectious contamination. There are numerous different kinds of growths. They reside on your skin, in the soil, on plants, in homes, and on household surfaces. Sometimes they can cause skin problems like rashes or bumps. The primary driver of contagious contamination is compromised resistance (either nearby resistance over the skin & mucous layers or foundational resistance as found on account of specific circumstances like diabetes & HIV/Helps).

Fig 1.5 Fungal Infection

1.6.1 Common Types

Some Common Types of Fungal Infection Include:

- Athlete's foot.
- Jock itch.
- Ringworm.
- Yeast infection.
- Onychomycosis, or a fungal infection of the nail [8].

1.7 Drugs & Excipients Profile

1.7.1 Bifonazole:

Source: Synthetic

Therapeutic Category: Azole antifungal drug.

Chemical Name: 1-[Phenyl (4-phenylphenyl) methyl]-1H-imidazole.

Chemical Formula: C22H18N2.

Molecular Structure:



Fig 1.6 M.S of Bifonazole

Physical & Chemical Properties:

Appearance: Solid.

M.W: 310.393g/mol

Colour: White to Pink or Salmon to Brown.

Purity: 99.9%.

Solubility: In water soluble

M.P: 145-151°C

Taste: Bitter.

Shape: Elliptical

MOA:

A compound called ergosterol, which is a crucial component of the parasitic cell layers, is produced by the body in a manner that biconazole inhibits. The parasite cytochrome p450 51 chemical, also known as lanosterol 14- alpha de-methylase, is affected by it. In cell memory design of growth, this is crucial. Cell lysis results from its restriction. Ergosterol's disruption of the cell film causes holes to become visible. The durability of organisms depends on their cell films. They prevent harmful elements from getting into phones and prevent contents from pouring out. Basic elements of parasitic cells may leak out as bifonazole causes openings to appear in cell films. It eliminates the organisms.

Side Effects:

- Dizziness.
- Nausea.
- Vomiting.
- Skin Rash.

- Redness of the Skin.
- Dry Skin.
- Vaginal Infection.

Contraindicated:

- It is contraindicated in patients oversensitive to the medication & other comparable anti-fungals.
- It ought to likewise not be utilized for nappy rash in children & during pregnancy & breast feeding in ladies.

Uses:

- It is used in the treatment of fungal infections like oral thrush, fungal infections of the skin such as jock itch, Athlete's foot.

Pharmacokinetic:

Absorption:
Following topical treatment, there was very little absorption (0.6% of the administered dose).

Half life: 1-2hrs.

Metabolism: Hepatic.

Density: .cm3/1g.±1.3

Storage: Room Temp.

1.7.2 Polymers Profile:

1.7.2.1 Propylene Glycol:

Synonyms: α-propylene, glycol, 1, 2-propanediol.

Molecular Structure:



Fig.1.7 M.S of Propylene Glycol

Molecular Formula: C3H5o2.

Chemical Name: Propane-1, 2- diol.

M.W: 76.09g/mol.

Anti-microbial Preservative, Disin-fectant & Humectants are functional categories.

Description:

A transparent, colourless substance that is viscous and practically odourless and has a sweet, slightly acerbic flavour, similar to glycerol.

Typical Characteristics

B.P: 188°C.

Flash Point: 99°C.

Freezing Point: -59°C.

Solubility:

Miscible with acetone, CCL4, 95% C2H5OH, glycerine & h2o.

Safety:

Propylene glycol is usually regarded as a moderately benign substance and is utilised in a wide range of medicinal compositions.

Storage & Stability:

Propylene glycol is stable at cool temperatures when tightly sealed, however at high temperatures when exposed to air, it will typically oxidise and produce things.

Applications:

- It is now widely used in a variety of parenteral & non-parenteral formulations as a dissolvable, extractant & additive.
- It dissolves a wide range of medications, including corticosteroids, phenols, sulfa medicines, barbiturates, nutrients (An & D), majority of alkaloids & many adjacent sedatives, making it a better general dissolvable than glycerine.
- As a germ-free substance, it is comparable to ethanol; against moulds, it is comparable to glycerine, but slightly less effective than ethanol [9].

1.7.2.2 Tween-20:

Synonyms: Armotan PML 20, Tween-20.

Chemical Structure:



Fig 1.8 Molecular Structure of Tween-20

Chemical Name: Poly-oxy-ethylene 20 sor-bitan mono-laurate.
Molecular Formula: C58H114O26.
M.W: 1128g/mol.
Functional Category: Nonionic surfactant; suspending Agent & Wetting Agent.
Description:
Colour: Yellow oily liquid at 25℃.
Odour & Taste: Poly-sorbates have a characteristic.
Applications:
Hydrophilic nonionic surfactants called poly-sorbates with 20 units of oxye-thylene are widely used as emulsifying agents in formulation of o/w medication emulsions.
Used as a wetting specialist in formulation of oral & perenteral suspensions; used as a solubilizing specialist for various compounds, such as natural ointments and nutrients that dissolve in oil.

Typical Properties:
Acid Value: 2.0%.
Specific Gravity: 1.1 at 25°C
Viscosity: 400mPas.
Surface Tension: 34.7mN/m at 20°C.
In-compatibilities:
Staining & precipitation happen with different substances, particularly phenols, tannins, tars & tarlike materials.
Stability & Storage Conditions:
Electrolytes, weak acids & weak bases do not affect poly-sorbates negatively; nevertheless, strong acids & strong bases cause gradual Saponification.
Poly-sorbates ought to be kept in a tightly closed, light-protected container that is kept in a cool, dry environment [37].

1.7.2.3 Carbopol-940:
Synonyms: 2-Propenoic acid Homopolymer, Acrylic acid Homopolymer.
Molecular Structure:



Fig.1.9 Molecular Structure of Carbopol-940

Molecular Formula: C18H34O2.
Chemical name: 2-octanoyloxypropyl Octanoate.
Non Proprietary Name BP, USP: Oleic acid.
Molecular Weight: 713.1g/mol.
Functional Category:
High viscosity polymer thickens carbo-mers effectively & produces sparkling, clear water or hydro-alcoholic gels.
Description:
Crosslinked Polyacrylic Acid Polymer is a white powder that creates dazzling clear gels or hydro-alcoholic gels & creams. It is a highly effective rheology modifier that can provide high viscosity.

Solubility: In water; after neutralization, they are soluble in 95% ethanol & glycerin.
Viscosity: 6.19mPas.
Specific Properties:
M.P: 116 °C.
Flash point: 130°F.
Moisture Content: 0.05 to 0.1 wt. %.
Surface tension: 24.3dynes/cm at 20°C.
Pharmaceutical Application:

- Carbopol 940 is much of the time utilized as a gelling specialist in gel arrangements.
- Grouping of Carbopol 940 as the need might arise to be worried to get a decent gel planning.
- This study expected to decide the impact of Carbopol 940 focus on actual properties, drug discharge and the utilization in eye drops.
- The examination strategy utilized was unmistakable technique with information assortment method utilizing PICO (Populace, Mediation, Think about, Outcame) approach.
- In light of Engaging exploration results, it was gotten that carbopol 940 had effects on the actual properties of the gel as pH, consistency, spreadability, bond, organoleptic & strength.
- Carbopol 940 is generally utilized in controlled-discharge drug definitions.

- Likewise, carbopol 940 is additionally protected to use in eye drops arrangements which don't cause disturbance in vivo testing.

Stability & Storage:
It needs to be kept in a small, tightly-sealed container in a cool, dry location away from light [10-12].

1.7.2.4 Methyl Paraben:
Nonproprietary Names:
BP: Methyl Hydroxy-benzente
USPNE: Methylparaben
Synonyms: 4-hydroxy benzoic acid & methyl p-hydroxy-benzoate
Chemical Name: Methyl-4-hydroxy-benzoate.
CAS Registry Number: 90-76-3.
Empirical Formula: $C_8H_8O_8$
Molecular Weight: 152.15
Functional Category:
Preser-vative as anti-microbial
Description:
Methylparaben occurs in the form of colorless or white crystalline form of powder. It is odorless & nearly odorless & a faint appearance having taste of burning sensation.
Melting point: 125-128℃.
Acidity/alkalinity: pH=4-8
Solubility
At 25^0 C solubility exist in H2O (2.5 g but considering benzene & CCl4.
Density: 1.352 g/cm^3
Stability:
Sterilized for 20 minutes by autoclave at 120℃. Deprived of de-composition.
Storage Conditions: Methyl Paraben should be stored in a well-closed jar.
Incompatibilities:
In the existence of non-ionic surfactants, such as poly-sorbate 80, anti-microbial activity of methyl-paraben & other parabens is considerably decreased as a consequence of micellisation.
Method of Manufacture:
By etherifying para-hydroxy-benzoic acid along with methanol, methyl-paraben is prepared.
Safety:
Preservatives as anti-microbial in cosmetics & oral & topical pharmaceutical formations methyl paraben & other parabens are extensively used. While in patenital & phthalmic preparations.
Healing Precautions:
Pay attention to standard measures necessary to conditions & quantity of material treated in a well-ventilated area it should be treated well, methyl-paraben may be irritating to the skin eyes.

1.7.2.6 Liquid Paraffin:

Synonyms:

Paraffinum liquidum, mineral oil, paraffinum perliquidum, paraffinum subliquidum.

Chemical or General Formula: $C2H_{2n+2}$

Functional Category:

Liquid paraffin is used as hydrating agent in various formulations such as hair shampoo & ointments to lock moisture into formulation.

Description:

Liquid paraffin all is a mineral oil that is a by product of the distillation of petroleum.

Boiling Point: 280-350^{0}C.

Viscosity: 7.5-100.0 cSt at 40^{0}C

Refractive Index: 1.476.

Solubility:

Practically insoluble, but soluble in organic solvents, ethanol (95percent), Glycerin & water Miscible, except fie castor oil, will volatile oils & with fixed oil.

Density: 0.877

Stability & Storage Conditions:

Responsible for oxidation during storage & light exposure, & stored in a cold, dry spot.

Incompatibilities:

Keep away from oxidizing agents & from highly acidic or alkaline materials.

Method of Manufacture:

Through distillation of petroleum, M ineral oil is Attained first, Distillation Cradicate the Hydrocarbon having lighter property & then redistillates the residue abut 330-390°C. **Safety:**

In a huge scope of drug details, mineral oil is used as an excipient mineral oil has been utilized remedially in constipation treatment since. When taken orally. It utilized to stool softening & as lubracants.

H&ling Precautions:

Pay attention to st&ard measures necessary to the conditions & quantity of material treated.

Regulatory Status:

Identified by GRAS Accepted or approved in the Unital Kingdom for usage in some applications for foodstuffs Encompassed in the FDA Database of Inactive ingredients.

1.7.2.7 Benzyl Alcohol:

Synonyms: Hydroxytoluol, Phenylcarbinol & Phenylmethanol.

Molecular Structure:



Fig 1.10 Molecular Structure of Benzyl Alcohol

Molecular Formula: C18H34O2.
Chemical name: Phenylmethanol.
Molecular Weight: 108.14g/mol.
Functional Category:
The overall dissolvable for inks, waxes, shellacs, veneers & epoxy gum coatings is benzyl liquor.
Description:

- It is a colourless liquid with a light, pleasant odour, and because of its polarity, low toxicity, low vapour pressure, it is an effective solvent.

Solubility:
It is miscible in CH3OH, diethyl ether & has a moderate solubility in H2O (4gm/ per 100millilitres).

Pharmacopeial Specifications:
Specific gravity: 1.045.
Viscosity: 5.474cPas.
Specific Properties:
M.P: -15°C.
Flash point: 93°C.
Surface tension: 72.8 dynes/cm.
Pharmaceutical Application:

- It is a sweet-smelling liquor that is used as a dissolvable, additive, scent fixing & consistency decreasing specialist in a large number of restorative details.
- A sweet-smelling corrosive called benzoic corrosive is used as an additive & pH agent in a great many restorative items.

Stability & Storage:
It needs to be kept in a small, tightly-sealed container that is cool, dry & away from light.

Chapter-2 Review of Literature

Venkatesh D.P.G, Karki R, et al., 2022 Bifonazole-stacked nanosponges were planned utilizing the emulsion dissolvable dispersion method. The nanosponges details were assessed with various boundaries. Molecule size & Zeta capability of Bifonazole stacked nanosponges for details were between the scope of 183.7 to 560.2nm & - 17.77 to - 21.9mV, individually. The surface morphology of nanosponges by SEM unveiled that it was round and permeable in nature. Drug entanglement productivity was viewed as 45.44 to 79.71%. The medication discharge study was finished by utilizing a phosphate cradle with a pH of 6.8. Further in vitro discharge information was fitted into active models. The upgraded definition M6 was consolidated with hydrogels to additional assessed utilizing boundaries, for example, in vitro drug discharge review, thickness, pH and skin aggravation concentrates on utilizing rodent model. Strength investigations of hydrogel definition MH2 uncovered no progressions in-vitro drug discharge, pH and medication content review toward the finish of a half year. In this way, the discoveries showed that the pre-arranged Bifonazole stacked nanosponges into hydrogel were steady. Subsequently, it very well may be a reasonable dose structure for the fix of parasitic diseases in the skin [13].

Alhakamy, N.A, et al. 2021 the results for luli-conazole-nano-emulgel (LUL-NEG) were 99.06 0.59%, 9.26 0.08 Pa.s, 5.65 0.17 & 1.31 0.08, respectively, for drug content examination, thickness, pH & refractive record. When compared to LUL gel, e saturation across rodent skin with LUL-NEG was thought to be overall higher. Additionally, skin irritation tests performed on exploratory species revealed that clear NEG and LUL-NEG did not produce any erythematous symptoms after 48 hours of exposure. Additionally, histo-pathological findings of exploratory skins did not reveal any peculiar symptoms at application site for strategy. In end, it was discovered that NEG definition created a truly crucial zone of obstruction (P 0.05) when compared to any other experimental groups. In conclusion, nano-emulgel method of delivering luli-conazole over skin to treat infectious skin illnesses may be a promising technique. [14].

Miastkowska et al., 2020 arranged nano-emulsion utilizing four medicinal ointments, including cinnamon rejuvenating balm. Point of study was to see fungicidal impact of President against parasitic strains. Results shows better fungicidal movement from arranged nanoemulsions utilizing medicinal balms than from unadulterated oils [15].

Botre P.P, Maniyar M.G, et al., 2020 Bifonazole is an imidazole antifungal medication utilized in type of salves. It was licensed in 1974 and endorsed for clinical use in 1983. Bifonazole having wide range action against dermatophytes, molds, yeasts, growths and a few gram positive microbes. BFZ SLNs frameworks were created by soften emulsification followed by dissolvable vanishing method utilizing Compritol 888ATO (Glyceryl behenate) as a strong lipid and Tween 80 as a surfactant. Created SLNs were assessed for molecule size, polydispersity list (PI), capture proficiency (EE) and medication discharge profiles. Process and definition boundaries were upgraded. Differential examining calorimetry (DSC) and X-

beam diffraction (XRD) studies were done on SLNs to check the progressions in the medication and lipid alterations. The BFZ SLNs based gels were arranged utilizing Carbopol 940 as a gelling specialist. The SLNs based gels were assessed for rheological boundaries, in vitro drug discharge and saturation studies. In vitro antifungal review recommended that the SLNs based gel was more successful in restraining development of C&ida albicans. Subsequently the review presumes that SLNs based gel of BFZ gives a supported delivery profile of BFZ and has the potential for treatment of skin parasitic contaminations [16].Chapter 3 Aim & Objectives

3.1 Aim:

Formulation & Evaluation of Bifonazole Nanoemulsion for the Treatment of Topical Fungal Infection

Objectives of the research work undertaken in detail are as follows:

- To conduct pre-formulation studies for the selection of oil & other co-formulating agents.
- To conduct compatibility studies of selected oil & co-formulating agents using API by FTIR to Formulate Bifonazole Nanoemulsion by using different surfactant mixture ratio.
- Characterization of the Formulation by Using Different Analytical Techniques.
- Designing of Antifungal Bifonazole Nanoemulsion of different concentration by using Polymers.
- In-vitro Evaluation of Formulated Bifonazole Nanoemulsion.
- To perform Microbiological Studies of Antifungal Cream & Comparison with Marketed Formulation.
- To Enhance Absorption of Cream by using suitable carrier incorporating in the formula.
- To Studied the physiochemical properties of prepared Bifonazole Nanoemulsion for an underst&ing of changes within these system during preparation & storage.

3.2 Plan of Work

3.2.1 Literature Review.

3.2.2 Procurement of Drugs.

3.2.3 Selection of Polymers.

3.2.4 Selection of Excipients for the preparation of nano-emulsion based formulation.

3.2.5 Preformulation Studies:

- Determination of Solubility of Drug in Different Solvent, Oil & Surfactants.
- Identification of Drug on the basis of Melting Point.
- Drug, Oil & Surfactant Compatibility Study by FTIR.
- Estimation of the Drug by UV Spectroscopy.
- Preparation of phase diagram.

3.2.6 Preparation of Nanoemulsions Formulation.

3.2.7 Evaluation of Formulation.

- Drug Content.
- Droplet size, PDI & Zeta Potential.
- Visual Inspection.
- Clarity test.
- Viscosity.
- pH.
- Spreadability.
- Homogeneity.
- % Yield.

3.2.8 Microbiological Studies.

3.2.9 Antifungal Activity Study by Using Cup Plate Method.

3.2.10 Skin Irritation Studies for Optimized Formulation.

3.2.11 Determination of In-vitro drug release of Nanoemulsions.

3.2.12 Stability Study.
3.2.13 Results & Discussion.
3.2.14 Conclusion.
3.2.15 References.

References:

1. Date AA, Naik B, Nagarsenker MS. Novel drug delivery systems: potential in improving topical delivery of antiacne agents. Skin Pharmacol Physiol 19(1):2- 16 (2006).
2. Schafer-Korting M, Korting HC, PoncePoschl E. Liposomal tretinoin for uncomplicated acne vulgaris. Clin Investig 72(12):1086-91 (1994 Dec).
3. Brisaert M, Grabriels M, Matthijs V, et al. Liposomes with tretinoin: a physical & chemical evaluation. J Pharm Biomed Anal 26(5-6):909-17 (2001 Dec).
4. Misra A.N.,"Controlled & Novel Drug Delivery", CBS Publishers & Distributors, New Delhi,1997, 107-109.
5. Shinoda, K., Carlsson, A. & Lindman, B., 1996. On the importance of hydroxyl groups in the polar head-group of nonionic surfactants & membrane lipids. *Advances in colloid & interface science*, *64*, pp.253-271.
6. Wynstra, F., Von Corswant, F. & Wetzels, M., 2010. In chains? An empirical study of antecedents of supplier product development activity in the automotive industry. *Journal of Product Innovation Management*, *27*(5), pp.625-639.
7. Aboofazeli, R., Lawrence, C.B., Wicks, S.R. & Lawrence, M.J., 1994. Investigations into the formation & characterization of phospholipid microemulsions. III. Pseudo-ternary phase diagrams of systems containing water-lecithin-isopropyl myristate & either an alkanoic acid, amine, alkanediol, polyethylene glycol alkyl ether or alcohol as cosurfactant. *International journal of pharmaceutics*, *111*(1), pp.63-72.
8. Pimpale, A., Pagare, R., Aher, S. & Phadtare, D., 2021. A Combine Spectrophotometric & Chromatographic Method Development & Validation of Levetiracetam Bulk & Tablet Dosage Form. *Asian Journal of Pharmaceutical Research & Development*, *9*(5), pp.40-49.
9. Lieberman HA, Rieger MM, Banker GS, Pharmaceutical dosage form: Disperse system, 2, Marcel Dekker, New York, 2005, 399-421.
10. Kikwai L, Babu RJ, Prado R, Kolot A, Armstrong CA, Ansel JC, Singh M, In-vitro & in-vivo evaluation of topical formulations of spantide II, AAPS Pharm Sci Tech, 6(4), 2005, 565-572.
11. Saroha K, Singh S, Aggarwal A, N&a S, Transdermal Gels- An Alternative Vehicle For Drug Delivery, Int Journal of Phr Chemical & Biological Sciences, 3(3), 2013, 495-503.

12. Aulton ME, Pharmaceutics: The Science of Dosage Form Ansel HC, Popovich NG Loyd VA, Pharmaceutical Dosage Forms & Drug Delivery Systems, 9, B. I. Publications, New Delhi, 2005, 407-408.
13. Gucwa, K., Milewski, S., Dymerski, T. & Szweda, P., 2018. Investigation of the antifungal activity & mode of action of Thymus vulgaris, Citrus limonum, Pelargonium graveolens, Cinnamomum cassia, Ocimum basilicum, & Eugenia caryophyllus essential oils. Molecules, 23(5), p.1116.
14. Maha E. Elmataeeshy a , Magda S. Sokar b , Mohammed Bahey-El-Din c , Dalia S. Shaker d, Enhanced transdermal permeability of Terbinafine through novel nanoemulgel formulation; Development, in vitro & in vivo characterization, Future Journal of Pharmaceutical Sciences xxx (2017) 1e11
15. Arvind Kaur *1 , Shaveta Sharma 1 , Mahavir Singh 2 , Peeyush Kaushik 1 , Abhilash Kutlehria, FORMULATION & EVALUATION OF NANO-EMULSION GEL OF AMPHOTERICIN B FOR TREATMENT OF SKIN INFECTIONS, Int. Res. J. Pharm. 2018, 9 (7)
16. Liu, X.H., Park, H., Hu, J.H., Hu, Y., Zhang, Q.L., Wang, B.L., Sun, B., Yeung, K.S., Zhang, F.L. & Yu, J.Q., 2017. Diverse ortho-C (sp2)–H functionalization of benzaldehydes using transient directing groups. Journal of the American Chemical Society, 139(2), pp.888-896.